

Student name _____

Treatment order date _____

Age _____ Weight _____

Treatment:

Diazepam rectal gel _____ mg rectally prn for:

- o seizure > _____ minutes--OR--for _____ or more seizures in _____ hours.

Use vagal nerve stimulator (VNS) magnet _____

Other _____

Call 911 if

- o Seizure does not stop by itself or with VNS within _____ minutes
- o Seizure does not stop within _____ minutes of giving diazepam rectal gel
- o Child does not start waking up within _____ minutes after seizure is over (no diazepam rectal gel given).
- o Child does not start waking up within _____ minutes after seizure is over (after diazepam rectal gel is given)

- Following a seizure: Child should rest in nurse's office
 Child may return to class
 Parents should be notified immediately
 Parents should receive a note/copy of the seizure record sent home with the child

Physician/Nurse Practitioner/Physician Assistant Name (Please print)

Signature/Date _____

License #/State _____

Address _____

Phone _____

Current medication(s) _____

Allergies _____

| Type of seizure(s) | Description* | |
|--|--|---|
| <input type="checkbox"/> Absence | <ul style="list-style-type: none">• Staring• Eye blinking | <ul style="list-style-type: none">• Loss of awareness• Other _____ |
| <input type="checkbox"/> Simple partial seizures | <ul style="list-style-type: none">• Remains conscious• Distorted sense of smell, hearing, sight | <ul style="list-style-type: none">• Involuntary rhythmic jerking/ twitching on one side• Other _____ |
| <input type="checkbox"/> Complex partial seizures | <ul style="list-style-type: none">• Confused• Not fully responsive/ unresponsive | <ul style="list-style-type: none">• May appear fearful• Purposeless, repetitive movements• Other _____ |
| <input type="checkbox"/> Generalized tonic-clonic seizures | <ul style="list-style-type: none">• Convulsions• Stiffening• Breathing may be shallow• Lips or skin may have bluish color | <ul style="list-style-type: none">• Unconsciousness• Confusion, weariness, or belligerence when seizure ends• Other _____ |

*Student may experience some or all of the listed symptoms during a specific seizure.

Seizure usually lasts _____ minutes and returns to baseline in _____ minutes.

Treatment for seizure _____

Call parents under the following circumstances:

1. _____

2. _____

Developed in collaboration with Christine O' Dell, RN, MSN, and Shlomo Shinnar,

MD, PhD, of the Comprehensive Epilepsy Management Center, Montefiore Medical Center, Bronx, New York

Figure 1. Seizure Emergency Treatment Plan